

JOURNAL OF MATHEMATICAL ANALYSIS AND APPLICATIONS **111**, 156–161 (1985)

On the Theorem of Power Expended in Continuum Mechanics

REMIGIO RUSSO

*Istituto di Matematica dell'Università,
via Mezzocannone, 8 – 80134 Naples, Italy*

Submitted by E. Stanley Lee

We show that the theorem of power expended may be claimed for unbounded bodies, under mild hypotheses on the behavior at infinity of the velocity, the density, and the stress fields. © 1985 Academic Press, Inc.

1. INTRODUCTION

Let E be an n -dimensional euclidean space, let B be a body, i.e., a regular region in E^1 , and let $\{\mathbf{0}, \mathbf{e}_i\}$ be an orthonormal reference frame in E . As known, the theorem of power expended assures that, if Ω is any bounded, fixed region contained at each instant on B , then

$$\begin{aligned} \frac{1}{2} \int_{\Omega} \rho v^2 d\Omega = \frac{1}{2} \int_{\Omega} \rho_0 v_0^2 d\Omega + \int_0^t ds \left\{ \int_{\Omega} \mathbf{b} \cdot \mathbf{v} d\Omega - \int_{\Omega} \mathbf{T} \cdot \mathbf{D} d\Omega \right. \\ \left. + \int_{\partial\Omega} \left(\mathbf{v} \cdot \mathbf{T} \mathbf{n} - \frac{1}{2} \rho v^2 \mathbf{v} \cdot \mathbf{n} \right) d\Sigma \right\} \end{aligned} \quad (1)$$

where the symbols have their usual meaning and the functions are supposed smooth [1].

This result poses the following question: *Does relation (1) hold if B and Ω are both unbounded, the body and surface forces are summable respectively on Ω and $\partial\Omega$, and the initial energy is finite?*

The following example [2] shows that such a generalization requires suitable restrictions upon the behavior at infinity of the flow $(\mathbf{v}, \rho, \mathbf{T})$. For homogeneous, incompressible, inviscid fluids $(\mathbf{T} = -p\mathbf{I})$, (1) takes the form

$$\frac{1}{2} \int_{\Omega} \rho v^2 d\Omega = \frac{1}{2} \int_{\Omega} \rho v_0^2 d\Omega + \int_0^t ds \left\{ \int_{\Omega} \mathbf{b} \cdot \mathbf{v} d\Omega - \int_{\partial\Omega} \left(p + \frac{1}{2} \rho v^2 \right) \mathbf{v} \cdot \mathbf{n} d\Sigma \right\}. \quad (2)$$

¹ Throughout this paper we follow the graphic conventions of [1], to which we refer for any symbols not explained here.

A simple consequence of (2) is the uniqueness of the null solution $(\mathbf{0}, \rho, -c\mathbf{I})$, with $c = \text{const}$, under the boundary condition $\mathbf{v} \cdot \mathbf{n} \equiv 0$. On the other hand, if $B \equiv E$ and $\mathbf{b} \equiv \mathbf{0}$, the initial boundary-value problem for the system of incompressible inviscid fluids admits the solution $(\sum_i t^2 \mathbf{e}_i, \rho, -2t \sum_i x_i \mathbf{I})$ as well as the trivial one. Hence, a necessary condition in order that (1) hold is $p = |\mathbf{T}| = o(r)$, with $r = |\mathbf{x} - \mathbf{0}|$.

A second restriction involves the velocity \mathbf{v} . Let $(\mathbf{v}, \rho, \mathbf{T})$ be a steady flow, let S_R be the sphere of radius R centered at $\mathbf{0}$, and let $\mathbf{v}_r(R) = \sup\{|\mathbf{v} \cdot \mathbf{e}_r|, \mathbf{x} \in S_R \cap \Omega, \mathbf{r}\mathbf{e}_r = \mathbf{x} - \mathbf{0}\}$. A physically reasonable request is that the integral $\int_0^{+\infty} dR/v_r$ is infinite.

Thus, as a first step, we may ask ourselves whether (1) holds for solutions $(\mathbf{v}, \rho, \mathbf{T})$ to the system

$$\begin{aligned} \rho \dot{\mathbf{v}} &= \text{div } \mathbf{T} + \mathbf{b} \\ \dot{\rho} + \text{div } \mathbf{v} &= 0 \end{aligned} \quad \text{on } Q = \Omega \times (0, +\infty) \quad (3)$$

under the hypotheses

$$\begin{aligned} &\text{there exists a positive and smooth function } q(r) \text{ on } [0, +\infty) \\ &\text{such that } q'(r) > 0, \lim_{r \rightarrow +\infty} q(r) = +\infty, q'(r) |\mathbf{v} \cdot \mathbf{e}_r| = O(1) \end{aligned} \quad (4)$$

$$\exists f(r) \in C((0, +\infty)), \quad f(r) = O(r), \quad \rho^{-1/2} |\mathbf{T}| = o(f(r)).$$

The aim of the present work is just to give a partly affirmative answer to the above question. In fact, denoting by Ξ the whole set $(\mathbf{v}, \rho, \mathbf{T})$ of solutions to system (3) such that²

$$\begin{aligned} \rho &\in C(\bar{Q}) \\ \mathbf{v} &\in C_1^1(Q) \cap C_1(\bar{Q}) \\ \mathbf{T} &\in C_2^1(Q) \cap C_2(\bar{Q}), \quad \mathbf{T} \cdot \mathbf{D} \geq 0 \end{aligned}$$

we shall prove the following

THEOREM. *Let B be an unbounded body, let $(\mathbf{v}, \rho, \mathbf{T}) \in \Xi$, and assume that*

² The set of functions defined on $G \subseteq E \times [0, +\infty)$ and differentiable up to the order m ($\in N_0$) inclusive will be denoted by $C^m(G)$ ($C^0(G) \equiv C(G)$) and we agree to use the symbol $C_k^m(G)$ for the set of tensor-valued functions of k th order, whose components belong to $C^m(G)$.

(4) is satisfied with $f^2(r) = r^{1-n}[q'(r)]^{-1}$. If Ω is any fixed region contained on B , $\forall t \geq 0$, and

$$\int_{\Omega} \rho_0 v_0^2 d\Omega < +\infty, \int_{\Omega} b^2/\rho d\Omega, \int_{\partial\Omega} |\mathbf{v} \cdot \mathbf{Tn} - \frac{1}{2}\rho v^2 \mathbf{v} \cdot \mathbf{n}| d\Sigma < +\infty, \forall t \geq 0$$

then (1) holds.

2. PROOF OF THE THEOREM

Consider the function

$$g(\mathbf{x}, s) = g_{\delta}(\psi(\mathbf{x}, t))$$

where g_{δ} is a C^{∞} function on \mathbb{R} verifying ($\delta > 0$)

$$\begin{aligned} g_{\delta}(\xi) &= 0 && \text{on } (-\infty, -\delta] \\ g_{\delta}(\xi) &= 1 && \text{on } [\delta, +\infty) \\ g'_{\delta}(\xi) &\geq 0 && \text{on } \mathbb{R} \end{aligned}$$

and

$$\psi(\mathbf{x}, s) = c^{-1}[q(R) + c(\tau - s) - q(r)], \quad R, \tau > 0.$$

The support of g , defined by the inequality

$$q(r) \leq q(R) + c(\tau - s),$$

is certainly compact on $E \times [0, \tau]$, $\forall R$. Moreover it is equal to the set

$$\chi_{\delta} = \{(\mathbf{x}, s) \in E \times [0, +\infty): r \leq q^{-1}[q(R) + c(\tau - s + \delta)], s \leq \tau\}.$$

Further, $g \in C^{\infty}(E \times [0, \tau])$ even if ψ is not differentiable at $\mathbf{0}$. Indeed, in the set

$$\{(\mathbf{x}, s) \in E \times [0, +\infty): r \leq q^{-1}[q(R) + c(\tau - s - \delta)], s \leq \tau\}$$

$g \equiv 1$. Hence, if we choose δ suitably small, we may conclude that g is differentiable at $\mathbf{0}$.

Multiply both sides of (3)₁ by $g\mathbf{v}$ and recall the following vector identities:

$$\begin{aligned} \frac{1}{2}g\rho\mathbf{v} \cdot \text{grad } v^2 &= \text{div} \left\{ \frac{1}{2}g\rho v^2 \mathbf{v} \right\} - \frac{1}{2}g v^2 \text{div}(\rho\mathbf{v}) - \frac{1}{2}\rho v^2 \mathbf{v} \cdot \text{grad } g \\ g\mathbf{v} \cdot \text{div } \mathbf{T} &= \text{div}\{g\mathbf{v} \cdot \mathbf{T}\} - g\mathbf{T} \cdot \mathbf{D} - \mathbf{v} \cdot \mathbf{T} \text{ grad } g. \end{aligned}$$

Integrate now over the set $\Omega_\mu \times (0, t)$, with $t < \tau$ and

$$\Omega_\mu = \{\mathbf{x} \in \Omega: |\mathbf{x} - \mathbf{y}| \geq \mu > 0, \forall \mathbf{y} \in \partial\Omega\},$$

and choose μ in such a way that $\Omega_\mu \neq \emptyset$.

A simple application of the divergence theorem leads at once to

$$\begin{aligned} \frac{1}{2} \int_{\Omega_\mu} g \rho v^2 d\Omega &= \frac{1}{2} \int_{\Omega_\mu} g_0 \rho_0 v_0^2 d\Omega - \frac{1}{2} \int_0^t ds \int_{\Omega_\mu} g'_\delta \rho v^2 d\Omega - \int_0^t ds \int_{\Omega_\mu} g \mathbf{T} \cdot \mathbf{D} d\Omega \\ &\quad - \frac{1}{2c} \int_0^t ds \int_{\Omega_\mu} g'_\delta q'(r) v^2 \mathbf{v} \cdot \mathbf{e}_r d\Omega \\ &\quad + \frac{1}{c} \int_0^t ds \int_{\Omega_\mu} g'_\delta q'(r) \mathbf{v} \cdot \mathbf{T} \mathbf{e}_r d\Omega \\ &\quad - \int_0^t ds \int_{\partial\Omega_\mu} g \left(\frac{1}{2} \rho v^2 \mathbf{v} - \mathbf{v} \cdot \mathbf{T} \right) \cdot \mathbf{n}_\mu d\Omega \end{aligned} \quad (5)$$

where \mathbf{n}_μ is the outward unit normal to $\partial\Omega_\mu$.

By virtue of the inequalities

$$\begin{aligned} -2 \frac{q'(r)}{c} \mathbf{v} \cdot \mathbf{T} \mathbf{e}_r &\leq \frac{1}{c} \rho v^2 + \frac{1}{c} [q'(r)]^2 T^2 / \rho \\ 2\mathbf{b} \cdot \mathbf{v} &\leq \rho v^2 + b^2 / \rho \end{aligned}$$

and since, by (4)₁, $\exists M > 0: q'(r) |\mathbf{v} \cdot \mathbf{e}_r| \leq M$ on $\Omega \times [0, \tau]$, (5) yields

$$\begin{aligned} \frac{1}{2} \int_{\Omega_\mu} g \rho v^2 d\Omega &\leq \frac{1}{2} \int_{\Omega_\mu} g_0 \rho_0 v_0^2 d\Omega + \frac{1}{2} \left(\frac{M+1}{c} - 1 \right) \int_0^t ds \int_{\Omega} g'_\delta \rho v^2 d\Omega \\ &\quad + \frac{1}{2} \int_0^t ds \int_{\Omega_\mu} g \rho v^2 d\Omega + \int_0^t ds \int_{\Omega_\mu} g (b^2 / 2\rho - \mathbf{T} \cdot \mathbf{D}) d\Omega \\ &\quad + \frac{1}{2c} \int_0^t ds \int_{\Omega_\mu} g'_\delta [q'(r)]^2 T^2 / \rho d\Omega \\ &\quad - \int_0^t ds \int_{\partial\Omega_\mu} g \left\{ \frac{1}{2} \rho v^2 \mathbf{v} - \mathbf{v} \cdot \mathbf{T} \right\} \cdot \mathbf{n}_\mu d\Sigma. \end{aligned} \quad (6)$$

Choose now $c > M + 1$. An obvious application of Grönwall's lemma and passage to the limit $\mu \rightarrow 0$ give

$$\begin{aligned} \frac{1}{2} \int_{\Omega} g \rho v^2 d\Omega &\leq e^t \left\{ \int_{\Omega} g_0 \rho_0 v_0^2 d\Omega + \int_0^t e^{-s} ds \left[\int_{\Omega} \left(b^2 / 2\rho - \mathbf{T} \cdot \mathbf{D} \right. \right. \right. \\ &\quad \left. \left. + \frac{1}{2c} g'_\delta [q'(r)]^2 T^2 / \rho \right) d\Omega - \int_{\partial\Omega} g \left(\frac{1}{2} \rho v^2 \mathbf{v} - \mathbf{v} \cdot \mathbf{T} \right) \cdot \mathbf{n} d\Sigma \right] \right\}. \end{aligned} \quad (7)$$

Observe now that, as $\delta \rightarrow 0$, g tends to the characteristic function of the set χ_0 , and the passage to the limit $\delta \rightarrow 0$ is permissible in (7). Moreover

$$\frac{1}{c} \int_0^t ds \int_{\Omega} g'_{\delta}[q'(r)]^2 T^2 / \rho d\Omega \xrightarrow{\delta \rightarrow 0} \int_0^t ds \int_{\Omega \cap \partial S_{R_s}} q'(r) T^2 / \rho d\Sigma$$

where we have put $R_s = q^{-1}[q(R) + c(\tau - s)]$. Thus, by letting $\mu \rightarrow 0$, (7) leads to

$$\begin{aligned} \frac{1}{2} \int_{\Omega \cap S_{R_t}} \rho v^2 d\Omega \leq e^t \left\{ \frac{1}{2} \int_{\Omega \cap S_{R_0}} \rho_0 v_0^2 d\Omega + \int_0^t e^{-s} ds \left[\int_{\Omega \cap S_{R_s}} (b^2/2\rho - \mathbf{T} \cdot \mathbf{D}) d\Omega \right. \right. \\ \left. \left. - \int_{\partial\Omega \cap S_{R_s}} (\frac{1}{2}\rho v^2 \mathbf{v} - \mathbf{v} \cdot \mathbf{T}) \cdot \mathbf{n} d\Sigma \right] + \int_0^t ds \int_{\Omega \cap \partial S_{R_s}} o(R^{1-n}) d\Sigma \right\}. \end{aligned}$$

This last relation assures us that ρv^2 and $\mathbf{T} \cdot \mathbf{D}$ are respectively summable on Ω and $\Omega \times (0, t)$, $\forall t \geq 0$. Indeed, letting $R \rightarrow +\infty$, we have

$$\begin{aligned} \frac{1}{2} \int_{\Omega} \rho v^2 d\Omega + \int_0^t ds \int_{\Omega} \mathbf{T} \cdot \mathbf{D} d\Omega \leq e^t \left\{ \frac{1}{2} \int_{\Omega} \rho_0 v_0^2 d\Omega + \int_0^t e^{-s} \left[\int_{\Omega} b^2/2\rho d\Omega \right. \right. \\ \left. \left. - \int_{\partial\Omega} (\frac{1}{2}\rho v^2 \mathbf{v} - \mathbf{v} \cdot \mathbf{T}) \cdot \mathbf{n} d\Sigma \right] \right\}. \quad (8) \end{aligned}$$

Now, let f be any smooth function of r . Multiplying both sides of (3)₁ by $f\mathbf{v}$ and operating in the same way as in the derivation of (5), we get, $\forall R > 0$,

$$\begin{aligned} \frac{1}{2} \int_{\Omega_{\mu} \cap S_R} f \rho v^2 d\Omega = \left\{ \frac{1}{2} \int_{\Omega_{\mu} \cap S_R} f \rho_0 v_0^2 d\Omega + \int_0^t ds \left[\int_{\Omega_{\mu} \cap S_R} (\frac{1}{2}\rho v^2 \mathbf{v} \text{ grad } f \right. \right. \\ \left. \left. - f \mathbf{T} \cdot \mathbf{D} - \mathbf{v} \cdot \mathbf{T} \text{ grad } f + \mathbf{b} \cdot \mathbf{v}) d\Omega \right. \right. \\ \left. \left. - \int_{\partial\Omega_{\mu} \cap S_R} f (\frac{1}{2}\rho v^2 \mathbf{v} - \mathbf{v} \cdot \mathbf{T}) \cdot \mathbf{n} d\Sigma \right. \right. \\ \left. \left. - \int_{\Omega_{\mu} \cap \partial S_R} f (\frac{1}{2}\rho v^2 \mathbf{v} - \mathbf{v} \cdot \mathbf{T}) \cdot \mathbf{e}_r d\Sigma \right] \right\}. \quad (9) \end{aligned}$$

By choosing $f \equiv 1$ in (9) and by virtue of the hypotheses and (8), we easily argue that the last integral, as $R \rightarrow +\infty$, is certainly finite. Then, let us choose $f = [q(r)]^{-\alpha}$, $\alpha > 0$. Since

$$\begin{aligned}
& - \int_{\Omega_\mu \cap S_R} \rho v^2 \mathbf{v} \operatorname{grad} f d\Omega \\
& = \alpha \int_{\Omega_\mu \cap S_R} [q(r)]^{-\alpha-1} q'(r) \rho v^2 \mathbf{v} \cdot \mathbf{e}_r d\Omega \\
& \leq K\alpha \int_{\Omega_\mu \cap S_R} \rho v^2 d\Omega \\
& - \int_{\Omega_\mu \cap S_R} \mathbf{v} \cdot \mathbf{T} \operatorname{grad} f d\Omega \\
& \leq \frac{1}{2}\alpha \int_{\Omega_\mu \cap S_R} \rho v^2 d\Omega \\
& + \frac{1}{2}\alpha \int d\gamma \int_0^R [q(r)]^{-2(\alpha+1)} [q'(r)]^2 T^2 / \rho r^{n-1} dr \\
& \leq \frac{1}{2}\alpha \int_{\Omega_\mu \cap S_R} \rho v^2 d\Omega + \frac{K\alpha}{2(2\alpha+1)} \{ [q(0)]^{-2\alpha-1} - [q(r)]^{-2\alpha-1} \}
\end{aligned}$$

where K is a suitable positive constant and (r, γ) is the spherical coordinate system associated to $\{\mathbf{0}, \mathbf{e}_i\}$, the theorem is finally achieved, by letting in (9) first $\alpha \rightarrow 0$, then $R \rightarrow +\infty$ and $\mu \rightarrow 0$.

3. CONCLUDING REMARKS

Our theorem may be easily proved for any unbounded material part P of B [1].

If we specify the material, we may obtain sharper results. A simple example is provided by an elastic flow, i.e., a fluid whose constitutive equation is $\mathbf{T} = -\pi(\rho) \mathbf{I}$. In this case assumption $(4)_2$ is no longer required. Moreover a very strong domain of influence theorem holds [3].

REFERENCES

1. M. E. GURTIN, "An Introduction to Continuum Mechanics," Academic Press, New York/London, 1981.
2. P. MURATORI, Teoremi di unicità per un problema relativo alle equazioni di Navier-Stokes, *Boll. Un. Mat. Ital.* (4) 4 (1971).
3. B. CARBONARO AND R. RUSSO, *About the system governing the motion of inviscid fluids*, submitted for publication.